

On Pinsker's Type Inequalities and Csiszár's f -divergences. Part I: Second and Fourth-Order Inequalities.

GUSTAVO L. GILARDONI *
Universidade de Brasília

Abstract

We study conditions on f under which an f -divergence D_f will satisfy $D_f \geq c_f V^2$ or $D_f \geq c_{2,f} V^2 + c_{4,f} V^4$, where V denotes variational distance and the coefficients c_f , $c_{2,f}$ and $c_{4,f}$ are *best possible*. As a consequence, we obtain lower bounds in terms of V for many well known distance and divergence measures. For instance, let $D_{(\alpha)}(P, Q) = [\alpha(\alpha - 1)]^{-1} [\int q^\alpha p^{1-\alpha} d\mu - 1]$ and $\mathcal{I}_\alpha(P, Q) = (\alpha - 1)^{-1} \log[\int p^\alpha q^{1-\alpha} d\mu]$ be respectively the *relative information of type* $(1 - \alpha)$ and *Rényi's information gain of order* α . We show that $D_{(\alpha)} \geq \frac{1}{2} V^2 + \frac{1}{72} (\alpha + 1)(2 - \alpha) V^4$ whenever $-1 \leq \alpha \leq 2$, $\alpha \neq 0, 1$ and that $\mathcal{I}_\alpha \geq \frac{\alpha}{2} V^2 + \frac{1}{36} \alpha(1 + 5\alpha - 5\alpha^2) V^4$ for $0 < \alpha < 1$. Pinsker's inequality $D \geq \frac{1}{2} V^2$ and its extension $D \geq \frac{1}{2} V^2 + \frac{1}{36} V^4$ are special cases of each one of these.

Keywords: Information or Kullback-Leibler divergence, relative entropy, variational or L^1 distance, Rényi's information gain, relative information.

MSC: 94A17, 26D15.

February 1, 2008

* *Address for correspondence:* Departamento de Estatística, Universidade de Brasília, Brasília, DF 70910-900, Brazil. E-mail: gilardon@unb.br. Tel/Fax: +5561 3273-6317

1 Introduction

Studying the relationship among the information divergence D and the variational distance V or, more specifically, determining lower bounds on D in terms of V , has been of interest at least since 1959, when Volkonskij and Rozanov [1] showed that $D \geq V - \log(1 + V)$. The best known result in this direction is usually referred to as Pinsker's inequality and states that $D \geq \frac{1}{2}V^2$. In general, studying the relationship between D and V is important because it allows one to "... translate results from information theory (results involving D) to results in probability theory (results involving V) and *vice versa*" (Fedotov, Harremoës and Topsøe, [2]). For instance, Barron [3] found a strengthened version of the central limit theorem by showing convergence in the sense of relative entropy and then using Pinsker's inequality to conclude convergence in the variational norm. In different settings, this idea has also been used by Topsøe [4] and Harremoës and Ruzankin [5]. Interestingly, these kind of results and its relation with Gagliardo-Nirenberg and generalized Sobolev inequalities have been used recently in order to obtain the decay rate of solutions of nonlinear diffusion equations—see Del Pino and Dolbeault [6] and references therein.

Pinsker's inequality was proved independently by Csiszár [7] and Kemperman [8], building on previous work by Pinsker [9], Kean [10] and Csiszár [11]. The constant $\frac{1}{2}$ in $D \geq \frac{1}{2}V^2$ is *best possible*, in the sense that there is a probability space and two sequences of probability measures P_n and Q_n such that $D(P_n, Q_n)/V^2(P_n, Q_n) \downarrow \frac{1}{2}$. Sharpened Pinsker type inequalities bounding D by higher-order polynomials in V^2 are also available. For instance, $D \geq \frac{1}{2}V^2 + \frac{1}{36}V^4$ —see Kullback [12, 13] and Vajda [14], where again the constant $\frac{1}{36}$ is best possible, in the sense that there are sequences P_n and Q_n such that $[D(P_n, Q_n) - \frac{1}{2}V^2(P_n, Q_n)]/V^4(P_n, Q_n) \downarrow \frac{1}{36}$. More recently, Topsøe showed in [15] that $D \geq \frac{1}{2}V^2 + \frac{1}{36}V^4 + \frac{1}{270}V^6 + \frac{221}{340200}V^8$, while Fedotov *et al* [2] have obtained a parametrization of the curve $v \mapsto L(v) = \inf\{D(P, Q) : V(P, Q) = v\}$ in terms of hyperbolic trigonometric functions and argue in [16] that the best possible extended Pinsker inequality contains terms up to and including V^{48} .

Let P and Q be probability measures on a measurable space $(\Omega, \mathcal{A})$ and p and q their densities or Radon-Nikodym derivatives with respect to a common dominating measure μ . The information divergence is $D(P, Q) = \int p \log(p/q) d\mu$ and is also known as *relative entropy* or *Kullback-Leibler divergence*. The variational (or L^1) distance is $V(P, Q) = \int |q - p| d\mu$. The f -divergence generated by f is $D_f(P, Q) = \int p f(q/p) d\mu$, where $f : (0, \infty) \rightarrow \mathbf{R}$ is convex and $f(1) = 0$. Jensen's inequality implies that $D_f(P, Q) \geq 0$ with equality holding if and only if $P = Q$, provided that f is strictly convex at $u = 1$. Hence, $D_f(P, Q)$ can be thought of as a measure of discrepancy between P and Q . The class of f -divergences was introduced by Csiszár [7] and Ali and Silvey [17]. It includes many of the most popular distances and discrepancy measures between probability measures. Both D and V belong to this class, respectively for $f(u) = -\log u$ and $f(u) = |u - 1|$. All of the following are also f -divergences: the χ^2 divergence $\chi^2(P, Q) = \int \frac{(q-p)^2}{p} d\mu = \int \frac{q^2}{p} d\mu - 1$, the Hellinger discrimination $h^2(P, Q) = \frac{1}{2} \int (\sqrt{q} - \sqrt{p})^2 d\mu = 1 - \int \sqrt{qp} d\mu$, the Triangular or Harmonic divergence $\Delta(P, Q) = \int \frac{(q-p)^2}{p+q} d\mu = 4 \int \frac{q^2}{p+q} d\mu - 2 = 2 - 4 \int \frac{pq}{p+q} d\mu$, the Capacitory discrimination $C(P, Q) = D(P, M) + D(Q, M)$ where $M = (P + Q)/2$ and the Jeffrey's divergence $J(P, Q) = D(P, Q) + D(Q, P) = \int (q - p) \log(q/p) d\mu$. A convenient one-parameter family which includes many of the above as special cases is generated by the convex functions $f(u) = [\alpha(\alpha - 1)]^{-1}(u^\alpha - 1)$ ($\alpha \neq 0, 1$). The resulting divergence $D_{(\alpha)}(P, Q) = [\alpha(\alpha - 1)]^{-1}[\int q^\alpha p^{1-\alpha} d\mu - 1]$ is called *relative information of type* $(1 - \alpha)$ by Vajda [18] and Taneja [19]. It is easy to check that $\chi^2 = 2D_{(2)}$, $4h^2 = D_{(1/2)}$ and $D = \lim_{\alpha \rightarrow 0} D_{(\alpha)}$. The Tsallis' and the Cressie-Read divergences, which are used

extensively in many areas including physics, economics and statistics, are respectively $T_\alpha = \alpha D_{(1-\alpha)}$ and $CR_\alpha = D_{(-\lambda)}$ (see [20, 21]). Finally, the Rényi's *information gain of order* $\alpha > 0$, $\mathcal{I}_\alpha(P, Q) = (\alpha - 1)^{-1} \log[\int p^\alpha q^{1-\alpha} d\mu]$, of which the information divergence is also a special case as $\alpha \rightarrow 1$, although not itself an f -divergence, can be expressed as $\mathcal{I}_\alpha = (\alpha - 1)^{-1} \log[1 - \alpha(1 - \alpha)D_{(1-\alpha)}]$, cf. [7, 20, 22].

Regarding the relationship between f -divergences and V , bounds are available for some special cases involving divergences which are more or less easy to manipulate. For instance, it is known that $\chi^2 \geq V^2$, $\frac{1}{2}V^2 \leq \Delta \leq V$ and $\frac{1}{8}V^2 \leq h^2 \leq \frac{1}{2}V$, see Dacunha-Castelle [23], LeCam [24], Dragomir, Glušćević and Pearce [25] and Topsøe [15]. A precise bound is available for the Capacitory divergence, for which Topsøe [26] showed that $C(P, Q) \geq \sum_{n=1}^{\infty} [n(2n - 1)2^{2n}]^{-1} V^{2n} = \log \frac{4-V^2}{4} + \frac{V}{2} \log \frac{2+V}{2-V}$.

Although we know of no general result giving a lower bound for f -divergences in terms of V , it appears as intuitively clear to us from the fact that f -divergences share many of the properties of D that inequalities similar to Pinsker's should also hold for other divergences. This should be the case, for instance, of relative information of type $(1 - \alpha)$ with α close to zero or that of Rényi's information gain of order α with α close to one. Maybe the closest to a general statement giving a kind of lower bound for an arbitrary D_f in terms of V is in Csiszár [27, Theorem 1], which states that $D_f(P, Q) < \epsilon$ implies, for sufficiently small ϵ , that $\frac{f''(1)}{2\sqrt{2}} V^2(P, Q) < \epsilon$ (cf. our Theorem 3 below), implying then that V should be small whenever D_f is small enough.

This paper will be the first of a series dealing with the relationship between f -divergences and variational distance. In particular, our objective here is to discuss conditions under which an f -divergence satisfies either a Pinsker's type inequality $D_f \geq c_f V^2$ or a fourth-order inequality $D_f \geq c_{2,f} V^2 + c_{4,f} V^4$. We will show in Section 3 that a sufficient condition for $D_f \geq c_f V^2$ is that the ratio between $(u - 1)^2$ and the difference between the generating f and its tangent at $u = 1$ be upper bounded by a straight line $a + bu$ with nonnegative coefficients a and b such that $a + b = c_f^{-1} = \frac{2}{f''(1)}$, if we want c_f to be *best possible*. A sufficient condition for having a fourth-order inequality, always with *best possible* coefficients, is presented in Section 4. Each of these theorems is followed by a corollary which gives conditions on the derivatives of f which are easier to check in practice than the original conditions on f . As a consequence of these we show in Section 3 that the relative information of type $(1 - \alpha)$ satisfies $D_{(\alpha)} \geq \frac{1}{2} V^2$ whenever $-1 \leq \alpha \leq 2$, $\alpha \neq 0, 1$. This inequality is improved in Section 4 to $D_{(\alpha)} \geq \frac{1}{2} V^2 + \frac{1}{72}(\alpha + 1)(2 - \alpha)V^4$. Using that $\mathcal{I}_\alpha = (\alpha - 1)^{-1} \log[1 - \alpha(1 - \alpha)D_{(1-\alpha)}]$, we also obtain that the Rényi's information gain of order α satisfies $\mathcal{I}_\alpha \geq \frac{1}{2} \alpha V^2 + \frac{1}{36} \alpha(1 + 5\alpha - 5\alpha^2) V^4$ whenever $0 < \alpha < 1$.

Besides sections 3 and 4 dealing respectively with second and fourth-order inequalities, the rest of the paper is organized as follows. Section 2 introduces some additional notation and states a fundamental inequality between powers of V and D_f in Corollary 2. In Section 5 we bring forward an argument from the sequel of this paper and briefly discuss why the tools that we use here to obtain second and fourth-order inequalities are insufficient to obtain sixth and higher-order inequalities when we are interested in *best possible* coefficients. Some technical results needed in Section 4 are presented in an Appendix. Finally, since some of the proofs in Section 4 and in the Appendix require somewhat lengthy calculations, we have recorded a MAPLE script that could help the reader interested in checking them. Although the script is not included here for reason of space, it is available from us on request. Notwithstanding, we stress that we have included in the paper what we believe are full and complete proofs for all statements made.

2 Notation and preliminary considerations

Throughout, equalities or inequalities between divergences will be understood to hold for every pair of probability measures, so that we will write for instance " $D \geq \frac{1}{2} V^2$ " instead of " $D(P, Q) \geq \frac{1}{2} V^2(P, Q)$ for every P, Q ". An equivalent definition for the Variational Distance is $V(P, Q) = 2 \sup\{|Q(A) - P(A)| : A \in \mathcal{A}\} = 2 [Q(B) - P(B)]$, where $B = \{\omega \in \Omega : q(\omega) \geq p(\omega)\}$. Hence $0 \leq V(P, Q) \leq 2$ with equality holding respectively if and only if $P = Q$ or $P \perp Q$. It is well known that the Information Divergence satisfies $0 \leq D(P, Q) \leq +\infty$. $D(P, Q) = 0$ can occur if and only if $P = Q$, while $P \not\ll Q$ implies that $D(P, Q) = +\infty$, although the reciprocal does not hold.

To avoid unnecessary discussion, we will assume the usual conventions $f(0) = \lim_{u \downarrow 0} f(u)$, $0 \cdot f(0/0) = 0$ and $0 \cdot f(a/0) = \lim_{\epsilon \downarrow 0} \epsilon f(a/\epsilon) = \lim_{u \uparrow +\infty} f(u)/u$. An f -divergence does not determine univocally the associated f . Indeed, for any a fixed, D_f and $D_{f-a(u-1)}$ are identical. For instance, $D = D_{-\log u} = D_{u-1-\log u}$. For any convex f we will let $\tilde{f}(u) = f(u) - f'(1)(u-1)$, which is nonnegative due to convexity considerations (more precisely, $f'(1)$ can be taken to be any number between the left and right derivatives of f at $u = 1$). We note here that second and higher-order derivatives of f and $\tilde{f}$ coincide. Indeed, we will often switch from one to the other in sections 3 and 4.

In general, f -divergences are not symmetric, in the sense that $D_f(P, Q)$ does not necessarily equals $D_f(Q, P)$, unless the generating f satisfies that $f(u) = uf(1/u) + a(u-1)$ for some fixed a . This is the case for instance of V and the h^2 and Δ divergences but not that of D or χ^2 . Whenever an f -divergence is not symmetric we could define the *reversed* divergence by letting $f_R(u) = uf(1/u)$, so that $D_{f_R}(P, Q) = D_f(Q, P)$. Similarly, beginning from an arbitrary f -divergence it is possible to construct a symmetric measure by using the convex function $f_S(u) = f(u) + f_R(u)$. For instance, the reversed information divergence is $D_R(P, Q) = \int q \log(q/p) d\mu$ and its symmetrized version is the already mentioned Jeffrey's divergence.

The following lemma is slightly more general than what we will actually need in Sections 3 and 4. It gives an upper bound on $|\int gq d\mu - \int gp d\mu| = |E_Q g - E_P g|$ in terms of a certain higher moment of g and an f -divergence D_f . If we interpret Q as an approximation to P , then $|E_Q g - E_P g|$ is the error between the corresponding approximate and actual expectations. The lemma generalizes results which we have used in [28, 29] in order to obtain upper limits for the approximating error in the context of Bayesian Statistics.

Lemma 1 *Let g be both P and Q integrable, $k(u) \geq 0$ such that $\int p k(q/p) d\mu = 1$, and $n > 1$. Then for any fixed a*

$$\left| \int gq d\mu - \int gp d\mu \right|^n \leq \sup_{u>0, u \neq 1} \left\{ \frac{(u-1)^n}{\tilde{f}(u)k^{n-1}(u)} \right\} \cdot [E_r |g-a|^{n/(n-1)}]^{n-1} \cdot D_f(P, Q), \quad (1)$$

where $r = p k(q/p)$, $E_r |g-a|^{n/(n-1)} = \int |g-a|^{n/(n-1)} r d\mu$ is the $[n/(n-1)]$ -th moment of g around a with respect to the probability density r and, as before, $\tilde{f}(u) = f(u) - f'(1)(u-1)$ and $f'(1)$ is a number between the left and the right derivative of f at $u = 1$.

PROOF: Let $m = n/(n-1)$ (so that n and m are conjugate) and $C = \{\omega : q(\omega) \neq p(\omega)\}$. Then we have for any real a that

$$\left| \int gq d\mu - \int gp d\mu \right| = \left| \int_C (g-a)(q-p) d\mu \right| \leq \int_C |g-a| |q-p| d\mu$$

$$\begin{aligned}
&= \int_C \left[\frac{|g-a| |q/p-1|}{\tilde{f}^{1/n}(q/p)} \right] \tilde{f}^{1/n}(q/p) p d\mu \leq \left[\int_C \frac{|g-a|^m |q/p-1|^m}{\tilde{f}^{m/n}(q/p)} p d\mu \right]^{1/m} \cdot [D_f(P, Q)]^{1/n} \\
&= \left[\int_C \frac{|g-a|^m |q/p-1|^m}{\tilde{f}^{m/n}(q/p) k(q/p)} k(q/p) p d\mu \right]^{1/m} \cdot [D_f(P, Q)]^{1/n} \\
&\leq \sup_{u>0, u \neq 1} \left\{ \frac{(u-1)^m}{\tilde{f}^{m/n}(u) k(u)} \right\}^{1/m} \cdot \left[\int |g-a|^m r d\mu \right]^{1/m} \cdot [D_f(P, Q)]^{1/n},
\end{aligned}$$

where the second inequality follows from Hölder's inequality. The desired result now follows after taking the n -th power in the leftmost and rightmost terms and noting that $\sup_{u>0, u \neq 1} \{(u-1)^m / \tilde{f}^{m/n}(u) k(u)\}^{n/m} = \sup_{u>0, u \neq 1} \{(u-1)^n / \tilde{f}(u) k^{n/m}(u)\} = \sup_{u>0, u \neq 1} \{(u-1)^n / \tilde{f}(u) k^{n-1}(u)\}$.

□

Taking $g = I_{p \geq q}$ and $a = \frac{1}{2}$, the left hand side of (1) becomes $2^{-n} |V(P, Q)|^n$, while $E_r |g-a|^{n/(n-1)} = \int |I_{p \geq q} - \frac{1}{2}|^{n/(n-1)} r d\mu \leq 2^{-n/(n-1)}$. Hence, we have the following corollary.

Corollary 2 *Let k , f and $\tilde{f}$ be as before. Then*

$$V^n(P, Q) \leq \sup_{u>0, u \neq 1} \left\{ \frac{(u-1)^n}{\tilde{f}(u) k^{n-1}(u)} \right\} \cdot D_f(P, Q) \quad (2)$$

REMARK 1. These results are still valid for nonconvex f provided that $\tilde{f}(u) \geq 0$ and we interpret $D_f(P, Q) = \int \tilde{f}(q/p) p d\mu$.

REMARK 2. Although (1) holds for any a , we usually would like to use a value for which $E_r |g-a|^{n/(n-1)}$ is small. Taking $a = E_r g = \int g r d\mu$ could be a good idea.

REMARK 3. A more precise formulation would use the L^∞ norm (or essential supremum) of $(u-1)^n / \tilde{f}(u) k^{n-1}(u)$ with respect to the measure $r d\mu$ instead of the supremum for $u > 0, u \neq 1$.

Of particular interest will be the cases $n = 2$ and $n = 4$, in which case equation (2) becomes

$$V^2(P, Q) \leq \sup_{u>0, u \neq 1} \left\{ \frac{(u-1)^2}{\tilde{f}(u) k(u)} \right\} D_f(P, Q) \quad (3)$$

and

$$V^4(P, Q) \leq \sup_{u>0, u \neq 1} \left\{ \frac{(u-1)^4}{\tilde{f}(u) k^3(u)} \right\} D_f(P, Q). \quad (4)$$

Some interesting inequalities follow directly from (3). For instance, taking (i) $k(u) = 1$ and $\tilde{f}(u) = (u-1)^2$ we obtain that $\chi^2 \geq V^2$, (ii) $k(u) = (1+u)/2$ and $\tilde{f}(u) = (u-1)^2/(1+u)$, so that $(u-1)^2/\tilde{f}(u)k(u) \equiv 2$, it follows that $\Delta \geq V^2$ and (iii) $k(u) = (\sqrt{u}+1)^2/[2(2-h^2(P, Q))]$ and $\tilde{f}(u) = \frac{1}{2}(\sqrt{u}-1)^2$, so that $(u-1)^2/\tilde{f}(u)k(u) \equiv 4[2-h^2(P, Q)]$, we obtain that $4h^2(2-h^2) \geq V^2$ —see Kraft [30], cited in Dragomir et al. [25]. Although we are not specially interested here in f -divergences for which $f''(1) = 0$, Corollary 1 also gives some bounds for this case. For instance, for the Triangular Divergence of order $\nu > 1$, $\Delta_\nu(P, Q) = \int \frac{(q-p)^{2\nu}}{(q+p)^{2\nu-1}} d\mu$ (see Topsøe [26] or Dragomir et al. [25]), we obtain after taking $n = 2\nu$, $\tilde{f}(u) = (u-1)^{2\nu}/(1+u)^{2\nu-1}$ and $k(u) = 1+u$ in (2) that $\Delta_\nu \geq V^{2\nu}$.




Figure 1: $h_w(u) = (u-1)^2 / (u-1-\log u)[w + (1-w)u]$ satisfies (i) $h_w(1) = 2$ for every w (by continuity) and (ii) when w is greater (smaller) than $\frac{1}{3}$, h_w attains its maximum for a u greater (respectively smaller) than 1, hence for $w \neq \frac{1}{3}$, the maximum value of h_w is greater than 2.

3 Second-order inequalities

We first note that also Pinsker's inequality follows from (3). To see this, take $\tilde{f}(u) = u - 1 - \log u$ and $k(u) = (1+2u)/3$ to obtain that $V^2 \leq \sup_{u>0, u \neq 1} h_{1/3}(u) \cdot D$, where $h_{1/3}(u) = 3(u-1)^2 / [(u-1-\log u)(1+2u)]$ (the reason for the subindex $1/3$ here will be made clear shortly). Now note that $\sup_{u>0, u \neq 1} h_{1/3}(u) = 2$ because $\lim_{u \rightarrow 1} h_{1/3}(u) = 2$, while $g(u) = 2(u-1-\log u)(1+2u) - 3(u-1)^2 \geq 0$ for $u > 0$ since $g(1) = g'(1) = 0$ and $g''(u) = 2(u-1)^2/u^2 \geq 0$. Observe that using $k(u) = (1+2u)/3$ in the previous argument amounts up to using the mixture $k(q/p)p = \frac{1}{3}p + \frac{2}{3}q$ in the proof of Lemma 1. It is interesting to note the reason why this and only this mixture works. Using a mixture $wp + (1-w)q$ for some $0 \leq w \leq 1$, $w \neq \frac{1}{3}$ is equivalent to taking $k(u) = 1 + (1-w)(1-u)$ in equation (3). Now, define $h_w(u) = (u-1)^2 / (u-1-\log u)[1 + (1-w)(u-1)]$ and observe that $\lim_{u \rightarrow 1} h_w(u) = 2$ for any w , so that $\sup_{u>0, u \neq 1} h_w(u) \geq 2$, while $h_w(u)$ attains its maximum value for $u = 1$ if and only if $w = \frac{1}{3}$. (see Figure 1). Hence, it follows that $\sup_{u>0, u \neq 1} h_w(u) > 2$ whenever $w \neq \frac{1}{3}$ and using any mixture with $w \neq \frac{1}{3}$ in Lemma 2 will produce a less than optimal inequality $D \geq cV^2$ with $c < \frac{1}{2}$.

The idea in the previous paragraph can be formulated for arbitrary f -divergences. Let $h_w(u) = (u-1)^2 / \{\tilde{f}(u)[1 + (1-w)(u-1)]\}$ and $b_w(u) = 1/h_w(u)$ and suppose that $f(u) = f'(1)(u-1) + \frac{1}{2}f''(1)(u-1)^2 + \frac{1}{6}f'''(1)(u-1)^3 + o(|u-1|^3)$ with $f''(1) \neq 0$. Equation (3) implies now that $D_f \geq [\sup_{u>0, u \neq 1} h_w(u)]^{-1}V^2$ for every $0 \leq w \leq 1$. Note that $\lim_{u \rightarrow 1} h_w(u) = 2/f''(1)$ for every w , so that $[\sup_{u>0, u \neq 1} h_w(u)]^{-1} \geq f''(1)/2$. In order to obtain the inequality $D_f \geq \frac{f''(1)}{2}V^2$, we must find a $w = w_f$ such that $[\sup_{u>0, u \neq 1} h_w(u)]^{-1} = f''(1)/2$. In other words, the (continuity corrected) function h_{w_f} should be maximized at $u = 1$, or equivalently $b_{w_f} = 1/h_{w_f}$ should be minimized at $u = 1$. It is easy to check that $b_w(u) = \frac{f''(1)}{2} + [\frac{f''(1)}{2}(1-w) + \frac{f'''(1)}{6}](u-1) + o(|u-1|)$. Hence, for b_{w_f}

to have a minimum at $u = 1$, the first-order term should vanish and hence $w_f = 1 + \frac{1}{3} \frac{f'''(1)}{f''(1)}$. Finally, for $h_{w_f}(u)$ to be actually maximized at $u = 1$, we must have that $h_{w_f}(u) \leq h_{w_f}(1) = 2/f''(1)$ and hence that

$$\tilde{f}(u)[1 + (1 - w_f)(u - 1)] \geq \frac{f''(1)}{2}(u - 1)^2. \quad (5)$$

This leads to the following theorem.

Theorem 3 (Pinsker type inequality for f -divergences). *Suppose that the convex function f is differentiable up to order 3 at $u = 1$ with $f''(1) > 0$, and let $w_f = 1 + \frac{1}{3} \frac{f'''(1)}{f''(1)}$. Then (5) implies that $D_f \geq \frac{f''(1)}{2} V^2$. The constant $\frac{f''(1)}{2}$ is best possible.*

PROOF: Although a rigorous proof can be obtained following the ideas above, the following argument is easier once the right condition has been identified. First, note that for (5) to hold we must have that $[1 + (1 - w_f)(u - 1)] > 0$ for every $u \neq 1$. Hence, (5) implies that

$$D_f(P, Q) = D_{\tilde{f}}(P, Q) \geq \frac{f''(1)}{2} \int \frac{(q/p - 1)^2}{1 + (1 - w_f)(q/p - 1)} p d\mu \geq \frac{f''(1)}{2} V^2(P, Q),$$

where the last inequality follows from (3) after taking $\tilde{f}(u) = (u - 1)^2/[1 + (1 - w_f)(u - 1)]$ and $k(u) = [1 + (1 - w_f)(u - 1)]$.

To show that c_f is best possible, consider a binary space and suppose that P assigns probabilities $p > 0$ and $(1 - p) > 0$ to each point of Ω , say $P = (p, 1 - p)$. For small v define $Q_v = (p + v/2, 1 - p - v/2)$ so that $V(P, Q_v) = v$ and $D_f(P, Q_v) = pf(1 + v/2p) + (1 - p)f(1 - v/2(1 - p)) = p \frac{f''(1)}{2} (v/2p)^2 + (1 - p) \frac{f''(1)}{2} [v/2(1 - p)]^2 + o(v^2) = \frac{f''(1)}{2} [4p(1 - p)]^{-1} v^2 + o(v^2)$. Hence $\lim_{v \rightarrow 0} D_f(P, Q_v)/V^2(P, Q_v) = \frac{f''(1)}{2} [4p(1 - p)]^{-1}$. Taking $p = \frac{1}{2}$ completes the proof. $\square$

REMARK. The last part of the proof suggests that, when there is no set A with $P(A) = \frac{1}{2}$, a better constant $c_f(P) = \frac{f''(1)}{2} [4 \sup\{P(A)[1 - P(A)] : A \in \mathcal{A}\}]^{-1} > \frac{f''(1)}{2}$ can be found so that $D_f(P, Q) \geq c_f(P) V^2(P, Q)$ for every Q . This problem has been addressed recently for the information divergence by Ordentlich and Weinberger [31].

For most divergences the condition in the next proposition is easier to check than (5).

Corollary 4 *Let f and $\tilde{f}$ be as before, $w_f = 1 + \frac{1}{3} \frac{f'''(1)}{f''(1)}$, and suppose that f is three times differentiable with $f''(u) > 0$ for all u . Then*

$$\text{sgn}(u - 1) \left\{ \frac{f'''(u)}{f''(u)} [1 + (1 - w_f)(u - 1)] + 3(1 - w_f) \right\} \geq 0 \quad (6)$$

implies (5) and hence that $D_f \geq \frac{f''(1)}{2} V^2$.

PROOF: Let $g(u) = \tilde{f}(u)[1 + (1 - w_f)(u - 1)] - \frac{f''(1)}{2}(u - 1)^2$. Now $g(1) = 0$, $g'(u) = \tilde{f}'(u)[1 + (1 - w_f)(u - 1)] + (1 - w_f)\tilde{f}(u) - f''(1)(u - 1)$ and hence $g'(1) = 0$, $g''(u) = \tilde{f}''(u)[1 + (1 - w_f)(u - 1)] + 2(1 - w_f)\tilde{f}'(u) - f''(1)$ and therefore $g''(1) = 0$ and finally $g'''(u) = f'''(u)[1 + (1 - w_f)(u - 1)] + 3(1 - w_f)f''(1)(u)$. Hence, (6) implies that $g'''(u) \leq 0$ for $u < 1$ and $g'''(1) \geq 0$ for $u > 1$, so the following Lemma implies that g must be nonnegative, which is equivalent to (5). $\square$

REMARK. Since we also have that $g'''(1) = 0$, (5) is also implied if $g^{(4)}(u) = f^{(4)}[1 + (1 - w_f)(u - 1)] + 4(1 - w_f)f'''(u) \geq 0$, but we usually find (6) easier to check.

Lemma 5 Let $n \geq 1$ and $g : (0, \infty) \rightarrow \mathbf{R}$ be $(n+1)$ times differentiable with $g(1) = g'(1) = \dots = g^{(n)}(1) = 0$, where $g^{(n)}$ is the n -th derivative of g , and suppose that either (i) n is even and $g^{(n+1)}(u) \leq 0$ for $u < 1$ and $g^{(n+1)}(u) \geq 0$ for $u > 1$ or (ii) n is odd and $g^{(n+1)}(u) \geq 0$ for every u . Then $g(u) \geq 0$ for every u .

PROOF: We prove first the case that n is odd. Since $g^{(n+1)}(u) \geq 0$ it follows that $g^{(n-1)}$ is convex, and since $g^{(n-1)}(1) = (g^{(n-1)})'(1) = 0$, it must have a minimum at $u = 1$, hence must be nonnegative. Repeat the argument backwards to obtain that $g^{(n-3)}, \dots, g$ are also nonnegative. Now in the case (i) that n is even, note that $g^{(n)}$ decreases for $u < 1$ and increases for $u > 1$ and, since $g^{(n)}(1) = 0$, it must be nonnegative, which reduces to the previous case. $\square$

In our view the most important application of Corollary 4 is to the relative information of type $(1 - \alpha)$ and the Rényi's information gain of order α .

Corollary 6 $D_{(\alpha)} \geq \frac{1}{2} V^2$ whenever $-1 \leq \alpha \leq 2$, $\alpha \neq 0, 1$. Also, $\mathcal{I}_\alpha \geq \frac{1}{2} \alpha V^2$ for $0 < \alpha < 1$. In both cases the coefficient of V^2 is best possible.

PROOF: For $D_{(\alpha)}$ we have $f(u) = [\alpha(\alpha - 1)]^{-1}(u^\alpha - 1)$. It is easy to check that $f''(1) = 1$, $w_f = (\alpha + 1)/3$ and the left hand side of (6) is $(\alpha + 1)(2 - \alpha)|u - 1|/3u$, which satisfies the condition for $-1 \leq \alpha \leq 2$.

Now, for $0 < \alpha < 1$, write $\mathcal{I}_\alpha = (1 - \alpha)^{-1} \log \left[1 + \frac{\alpha(1 - \alpha)D_{(1 - \alpha)}}{1 - \alpha(1 - \alpha)D_{(1 - \alpha)}} \right]$ and use that $\log(1 + x) \geq \frac{2x}{2 + x}$ for $x \geq 0$ (cf. Topsøe [32]) to get that $\mathcal{I}_\alpha \geq \alpha D_{(1 - \alpha)} \frac{2}{2 - \alpha(1 - \alpha)D_{(1 - \alpha)}} \geq \frac{1}{2} \alpha D_{(1 - \alpha)}$. $\square$

REMARK 1. Pinsker's inequality can be seen as the limiting case of the inequality just stated for $D_{(\alpha)}$ as $\alpha \rightarrow 0$ or equivalently of that stated for $\mathcal{I}_\alpha$ as $\alpha \uparrow 1$.

REMARK 2. The behavior of $\mathcal{I}_\alpha$ for $\alpha > 1$ is somewhat puzzling to us. We will show in Section 4 (cf. Theorem 7 and Corollary 9) that for any $-1 \leq \alpha \leq 2$ there are probability measures P_v and Q_v such that $V(P_v, Q_v) = v$ and $D_{(1 - \alpha)}(P_v, Q_v) = \frac{1}{2} v^2 + \frac{1}{72} (\alpha + 1)(2 - \alpha) v^4 + o(v^4)$. Hence, we must have that $\mathcal{I}_\alpha(P_v, Q_v) = (\alpha - 1)^{-1} \log[1 - \alpha(1 - \alpha)D_{(1 - \alpha)}(P_v, Q_v)] = \frac{1}{2} \alpha v^2 + \frac{1}{36} \alpha(1 + 5\alpha - 5\alpha^2) v^4 + o(v^4)$. Since $1 + 5\alpha - 5\alpha^2 < 0$ for $\alpha > \frac{1}{2} + \frac{3}{10} \sqrt{5} \approx 1.17$, it follows that in this case we cannot have that $\mathcal{I}_\alpha \geq \frac{1}{2} \alpha V^2$. We do not know whether the inequality holds for $1 < \alpha \leq \frac{1}{2} + \frac{3}{10} \sqrt{5}$.

4 Fourth-order inequalities

The inequality $D \geq \frac{1}{2} V^2 + \frac{1}{36} V^4$ is a consequence of the fact that

$$u - 1 - \log u \geq \frac{1}{2} \frac{(u - 1)^2}{1 + \frac{2}{3}(u - 1)} + \frac{1}{36} \frac{(u - 1)^4}{[1 + \frac{28}{45}(u - 1)]^3} \quad (7)$$

for all $u > 0$ together with (3) and (4). To prove (7), let $g(u) = (u - 1 - \log u)[1 + \frac{2}{3}(u - 1)][1 + \frac{28}{45}(u - 1)]^3 - \frac{1}{2}(u - 1)^2[1 + \frac{28}{45}(u - 1)]^3 - \frac{1}{36}(u - 1)^4[1 + \frac{2}{3}(u - 1)]$ and use Lemma 5 after showing that $g(1) = g'(1) = g''(1) = g'''(1) = g^{(4)}(1) = g^{(5)}(1) = 0$ and $g^{(6)}(u) = \frac{8}{91125}(43904u^4 - 50960u^3 + 44268u^2 - 34102u + 24565)/u^6$ is positive everywhere because

$$\begin{aligned} & 43904u^4 - 50960u^3 + 44268u^2 - 34102u + 24565 \\ &= 43904(u - \frac{65}{224})^4 + \frac{88347}{4}(u - \frac{1907903}{2827104})^2 + \frac{10273158845617}{723738624}. \end{aligned} \quad (8)$$

In this section we generalize the idea in the last paragraph for arbitrary f -divergences. In other words, an inequality of the form $D_f \geq c_{2,f}V^2 + c_{4,f}V^4$ would be obtained if we can prove that

$$\tilde{f}(u) \geq c_{2,f} \frac{(u-1)^2}{1 + (1 - w_{2,f})(u-1)} + c_{4,f} \frac{(u-1)^4}{[1 + (1 - w_{4,f})(u-1)]^3}. \quad (9)$$

for all $u > 0$. For this inequality to hold and being sharp enough so that $c_{2,f}$ and $c_{4,f}$ are best possible, it is necessary that the Taylor expansions of both sides around $u = 1$ must coincide up to and including fifth-order terms. This condition implies the expression of the $c_{i,f}$'s and $w_{i,f}$'s ($i = 2, 4$) in terms of the derivatives of f at $u = 1$ in the next theorem.

Theorem 7 (Fourth-order extended Pinsker inequality for f -divergences). *Let f and $\tilde{f}$ be as before and define $c_{2,f} = \frac{f''(1)}{2}$, $w_{2,f} = 1 + \frac{1}{3} \frac{f'''(1)}{f''(1)}$, $c_{4,f} = \frac{1}{72} [3f^{(4)}(1) - 4 \frac{(f''')^2(1)}{f''(1)}]$ and*

$$w_{4,f} = 1 + \frac{1}{45} \frac{9f^{(5)}(1) - 20 \frac{(f''')^3(1)}{(f'')^2(1)}}{3f^{(4)}(1) - 4 \frac{(f''')^2(1)}{f''(1)}}.$$

Suppose that both $c_{2,f} > 0$ and $c_{4,f} > 0$ and that for every $u > 0$ we have

$$\begin{aligned} & \tilde{f}(u) [1 + (1 - w_{2,f})(u-1)] [1 + (1 - w_{4,f})(u-1)]^3 \\ & \geq c_{2,f} (u-1)^2 [1 + (1 - w_{4,f})(u-1)]^3 + c_{4,f} (u-1)^4 [1 + (1 - w_{2,f})(u-1)]. \end{aligned} \quad (10)$$

Then $D_f \geq c_{2,f}V^2 + c_{4,f}V^4$. The coefficient $c_{4,f}$ is best possible.

PROOF: We will prove first that (10) implies that $0 \leq w_{i,f} \leq 1$ ($i = 2, 4$). Reasoning by contradiction, suppose first that $w_{2,f} \notin [0, 1]$, and evaluate both sides of (10) at $u = -w_{2,f}/(1 - w_{2,f}) > 0$ to obtain that $0 \geq c_{2,f}(1 - w_{2,f})^5(w_{4,f} - w_{2,f})^3$. Since $c_{2,f} > 0$, this implies that $w_{4,f} \geq w_{2,f}$ when $w_{2,f} > 1$ and $w_{4,f} \leq w_{2,f}$ when $w_{2,f} < 0$. Hence also $w_{4,f} \notin [0, 1]$ and we can evaluate again both sides of (10) now at $u = -w_{4,f}/(1 - w_{4,f}) > 0$ to get that $0 \geq c_{4,f}(1 - w_{4,f})^5(w_{2,f} - w_{4,f})$, so that $w_{4,f} > 1$ implies now that $w_{2,f} \geq w_{4,f}$ while $w_{4,f} < 0$ implies that $w_{2,f} \leq w_{4,f}$. Hence we must have in any case that $w_{2,f}$ and $w_{4,f}$ are equal, say $w_{2,f} = w_{4,f} = w$, so that (10) becomes

$$\tilde{f}(u) [1 + (1 - w)(u-1)]^4 \geq c_{2,f} (u-1)^2 [1 + (1 - w)(u-1)]^3 + c_{4,f} (u-1)^4 [1 + (1 - w)(u-1)]. \quad (11)$$

Since we are still assuming that $w_{2,f} = w \notin [0, 1]$, consider this inequality as $u \rightarrow -w/(1 - w) > 0$. The left hand side is equivalent to $f[-w/(1 - w)](1 - w)^4[u + w/(1 - w)]^4$, i.e. a positive coefficient times an infinitesimal term of order 4 in $[u + w/(1 - w)]$, while the right hand side is equivalent to $c_{4,f}(1 - w)^3[u + w/(1 - w)]$, i.e. an infinitesimal term of order 1 in $[u + w/(1 - w)]$. If $w > 1$ we have that the principal part here is $c_{4,f}(1 - w)^3 < 0$ and hence (11) cannot hold as $u \uparrow -w/(1 - w)$, while if $w < 0$ the principal part of the right hand side is $c_{4,f}(1 - w)^3 > 0$ and (11) cannot hold as $u \downarrow -w/(1 - w)$. This contradiction is due to the assumption that $w_{2,f} \notin [0, 1]$. Similarly we prove that $0 \leq w_{4,f} \leq 1$.

Now that we have proved that both $[1 + (1 - w_{2,f})(u-1)]$ and $[1 + (1 - w_{4,f})(u-1)]$ are positive for $u > 0$, we observe that Condition (10) implies (9) and hence that

$$\begin{aligned} D_f(P, Q) &= D_{\tilde{f}}(P, Q) \\ &\geq c_{2,f} \int \frac{(q/p - 1)^2}{1 + (1 - w_{2,f})(q/p - 1)} p d\mu + c_{4,f} \int \frac{(q/p - 1)^4}{[1 + (1 - w_{4,f})(q/p - 1)]^3} p d\mu. \end{aligned}$$

Hence, use respectively (3) and (4) to bound each term in the right hand side to obtain that $D_f \geq c_{2,f} V^2 + c_{4,f} V^4$.

The proof that $c_{4,f}$ is best possible is similar to the last part of the proof of Theorem 3. Consider again a binary space and for small v define $P = (p, 1 - p)$ and $Q_v = (p + v/2, 1 - p - v/2)$ so that $V(P, Q_v) = v$ and $D_f(P, Q_v) = pf(1 + v/2p) + (1 - p)f(1 - v/2(1 - p))$. Now let $p = \frac{1}{2} + \frac{1}{6} \frac{f'''(1)}{f''(1)} v$ and expand $D_f(P, Q_v)$ around $v = 0$ to obtain that $D_f(P, Q) = c_{2,f} v^2 + c_{4,f} v^4 + o(v^4)$. We leave the details to the reader. $\square$

The following condition on the derivatives of f is usually easier to prove than (10).

Corollary 8 *Let $f, \tilde{f}, c_{2,f}, w_{2,f}, c_{4,f}$ and $w_{4,f}$ be as before, and suppose that f is six times differentiable with $f''(u) > 0$ for all u . Then*

$$\begin{aligned} & \frac{f^{(6)}(u)}{f''(u)} [1 + (1 - w_{2,f})(u - 1)][1 + (1 - w_{4,f})(u - 1)]^3 \\ & + 6 \frac{f^{(5)}(u)}{f''(u)} [1 + (1 - w_{4,f})(u - 1)]^2 [4 - w_{2,f} - 3w_{4,f} + 4(1 - w_{2,f})(1 - w_{4,f})(u - 1)] \\ & + 90 \frac{f^{(4)}(u)}{f''(u)} (1 - w_{4,f}) [1 + (1 - w_{4,f})(u - 1)] [2 - w_{2,f} - w_{4,f} + 2(1 - w_{2,f})(1 - w_{4,f})(u - 1)] \\ & + 120 \frac{f'''(u)}{f''(u)} (1 - w_{4,f})^2 [4 - 3w_{2,f} - w_{4,f} + 4(1 - w_{2,f})(1 - w_{4,f})(u - 1)] \\ & + 144(1 - w_{2,f})(1 - w_{4,f})^3 \geq 0 \end{aligned} \tag{12}$$

implies (10) and hence that $D_f \geq c_{2,f} V^2 + c_{4,f} V^4$.

PROOF: Let $g(u) = \tilde{f}(u)[1 + (1 - w_{2,f})(u - 1)][1 + (1 - w_{4,f})(u - 1)]^3 - c_{2,f}(u - 1)^2[1 + (1 - w_{4,f})(u - 1)]^3 - c_{4,f}(u - 1)^4[1 + (1 - w_{2,f})(u - 1)]$. Now it is straightforward, although rather tedious, that $g(1) = g'(1) = g''(1) = g'''(1) = g^{(4)}(1) = g^{(5)}(1) = 0$, while $g^{(6)}(u)$ equals the left hand side of (12) times $f''(u)$ and hence is nonnegative. Hence, Lemma 5 implies that $g(u) \geq 0$, which is equivalent to (10). $\square$

The case of Jeffrey's divergence ($f(u) = \tilde{f}(u) = (u - 1) \log u$) is interesting, because we have that $c_{2,f} = 1$, $w_{2,f} = \frac{1}{2}$, $c_{4,f} = \frac{1}{12}$, $w_{4,f} = \frac{1}{2}$ and the left hand side of (12) is $\frac{3}{2}(5u^4 - 8u^3 + 9u^2 - 8u + 5)/u^4 = \frac{3}{2}[5(u - \frac{2}{5})^4 + \frac{21}{5}(u - \frac{4}{5})^2 + \frac{273}{125}]/u^4 > 0$. Hence $J \geq V^2 + \frac{1}{12}V^4$. We note that from $D \geq \frac{1}{2}V^2 + \frac{1}{36}V^4$ it follows immediately using the symmetry of V that $J(P, Q) = D(P, Q) + D(Q, P) \geq V^2 + \frac{1}{18}V^4$, but this bound is worst than the one we found using Corollary 8. To finish this section we present the special case of the relative information and information gain of order α .

Corollary 9 $D_{(\alpha)} \geq \frac{1}{2}V^2 + \frac{1}{72}(\alpha + 1)(2 - \alpha)V^4$ for $-1 \leq \alpha \leq 2$, $\alpha \neq 0, 1$. Also, $\mathcal{I}_\alpha \geq \frac{\alpha}{2}V^2 + \frac{1}{36}\alpha(1 + 5\alpha - 5\alpha^2)V^4$ for $0 < \alpha < 1$. In both cases the coefficients of V^4 are best possible.

PROOF: We prove first the assertion for $D_{(\alpha)}$. We have $f(u) = [\alpha(\alpha - 1)]^{-1}(u^\alpha - 1)$ and then $c_{2,f} = \frac{1}{2}$, $w_{2,f} = \frac{\alpha+1}{3}$, $c_{4,f} = \frac{1}{72}(\alpha + 1)(2 - \alpha)$, $w_{4,f} = \frac{17+11\alpha}{45}$. Again straightforwardly although rather lengthy, the left hand side of (12) is

$$\frac{(\alpha + 1)(2 - \alpha)}{273375 u^{-4}} [-(\alpha - 5)(\alpha - 3)(\alpha - 4)(11\alpha + 17)^3]$$

$$\begin{aligned}
& +2(\alpha-3)(\alpha-4)(22\alpha^2-28\alpha-59)(11\alpha+17)^2u \\
& -6(11\alpha+17)(-28+11\alpha)(\alpha-3)(\alpha+2)(11\alpha^2-11\alpha-31)u^2 \\
& +2(3+\alpha)(\alpha+2)(22\alpha^2-16\alpha-65)(-28+11\alpha)^2u^3 \\
& -(\alpha+4)(3+\alpha)(\alpha+2)(-28+11\alpha)^3u^4 \Big], \tag{13}
\end{aligned}$$

which we prove in the appendix that is positive for $u > 0$ when $-1 \leq \alpha \leq 2$.

Now, for Rényi's information gain we proceed as in the proof of Corollary 4 and use again that $\log(1+x) \geq \frac{2x}{2+x}$ for $x \geq 0$, so that

$$\begin{aligned}
\mathcal{I}_\alpha &= (1-\alpha)^{-1} \log \left[1 + \frac{\alpha(1-\alpha)D_{(1-\alpha)}}{1-\alpha(1-\alpha)D_{(1-\alpha)}} \right] \\
&\geq \alpha D_{(1-\alpha)} \frac{2}{2-\alpha(1-\alpha)D_{(1-\alpha)}} \geq \alpha D_{(1-\alpha)} \left[1 + \frac{1}{2}\alpha(1-\alpha)D_{(1-\alpha)} \right] \\
&\geq \frac{1}{2}V^2 + \frac{1}{72}(\alpha+1)(2-\alpha)V^4 + \frac{\alpha^2}{2}(1-\alpha)\frac{1}{4}V^4 = \frac{\alpha}{2}V^2 + \frac{1}{36}\alpha(1+5\alpha-5\alpha^2)V^4.
\end{aligned}$$

□

Using Corollary 9 we can obtain bounds for the Tsallis and Cressie-Read divergences, since they can be expressed in terms of $D_{(\alpha)}$.

5 A note about higher-order inequalities

Unfortunately, the tools we have used so far are insufficient to obtain inequalities of the form $D_f \geq c_{2,f}V^2 + c_{4,f}V^4 + \dots + c_{2n,f}V^{2n}$ including terms of at least order 6. To understand why, it is useful to restrict attention to the information divergence D . Our discussion at the beginning of Sections 2 and 3 shows that the second and fourth-order inequalities for D are due respectively to the inequalities $u-1-\log u \geq \frac{1}{2}(u-1)^2/[1+\frac{2}{3}(u-1)]$ and (7) and Corollary 2. Now, it is straightforward to check that the difference between the left and the right hand sides of (7) equals $\frac{41}{12150}(u-1)^6 + O(|u-1|^7)$. Indeed, we conjecture that

$$u-1-\log u \geq \frac{1}{2} \frac{(u-1)^2}{1+\frac{2}{3}(u-1)} + \frac{1}{36} \frac{(u-1)^4}{[1+\frac{28}{45}(u-1)]^3} + \frac{41}{12150} \frac{(u-1)^6}{[1+\frac{23186}{38745}(u-1)]^5}. \tag{14}$$

However, even if we could prove this assertion, we would obtain then only that $D \geq \frac{1}{2}V^2 + \frac{1}{36}V^4 + \frac{41}{12150}V^6$. The coefficient $\frac{41}{12150}$, even if close, is smaller than the best possible $\frac{1}{270}$ found by Topsøe [15]. A possible explanation for this, or in other words, from where the difference $\frac{1}{270} - \frac{41}{12150} = \frac{2}{6075}$ comes from, can be obtained looking at the divergences which result from the right hand side of (14). We have on one side that $\int \frac{(q/p-1)^2}{[1+(2/3)(q/p-1)]^3} p d\mu \geq V^2(P, Q)$, with equality holding if and only if $[1+\frac{2}{3}(q/p-1)] \propto |q/p-1|$ (actually, the proportionality constant must equal $1/V(P, Q)$). Similarly, $\int \frac{(q/p-1)^4}{[1+(28/45)(q/p-1)]^3} p d\mu \geq V^4(P, Q)$, with equality holding if and only if $[1+\frac{28}{45}(q/p-1)] \propto |q/p-1|$. Hence, it follows from these two inequalities that

$$\frac{1}{2} \int \frac{(q/p-1)^2}{[1+(2/3)(q/p-1)]^3} p d\mu + \frac{1}{36} \int \frac{(q/p-1)^4}{[1+(28/45)(q/p-1)]^3} p d\mu \geq \frac{1}{2}V^2(P, Q) + \frac{1}{36}V^4(P, Q), \tag{15}$$

but now equality cannot hold since, even if close, $\frac{2}{3} \neq \frac{28}{45}$. In fact, we conjecture that the infimum of the left hand side of (15) taken over all P and Q such that $V(P, Q) = v$ equals $\frac{1}{2}v^2 + \frac{1}{36}v^4 + (\frac{1}{270} - \frac{41}{12150})v^6 + o(v^6)$. We hope to be able to report on these issues soon.

6 Appendix

Before proving (13) we will state the following lemma. We have already used the idea in the lemma to obtain the decomposition (8).

Lemma 10 *Let $T(u) = c_4u^4 + c_3u^3 + c_2u^2 + c_1u + c_0$ and define $a_4 = c_4$, $a_2 = \frac{1}{8}[8c_2c_4 - 3c_3^2]/c_4$ and*

$$a_0 = \frac{1}{256c_4^3} \frac{2048c_0c_4^4c_2 - 768c_0c_4^3c_3^2 - 8c_3^4c_2c_4 + c_3^6 + 64c_3^3c_1c_4^2 - 512c_1^2c_4^4}{8c_2c_4 - 3c_3^2}. \quad (16)$$

Then a sufficient condition for $T(u) \geq 0$ for every u is that a_4 , a_2 and a_0 are nonnegative.

PROOF: Check that

$$T(u) = a_4 \left(u + \frac{c_3}{4c_4}\right)^4 + a_2 \left(u + \frac{1}{4c_4} \frac{16c_1c_4^2 - c_3^3}{8c_2c_4 - 3c_3^2}\right)^2 + a_0.$$

PROOF OF (13): Let $-1 \leq \alpha \leq 2$, $T(u) = T_\alpha(u)$ be the term between brackets in (13) and define $c_i = c_i(\alpha)$ and $a_i = a_i(\alpha)$ as in Lemma 10, so that for instance $a_4 = c_4 = -(\alpha + 4)(\alpha + 3)(\alpha + 2)(11\alpha - 28)^3$ which is of course nonnegative. Next □

$$a_2 = \frac{9}{2} \frac{(980\alpha^3 + 552\alpha^2 - 4257\alpha - 4207)(\alpha + 2)(-28 + 11\alpha)}{\alpha + 4},$$

which is positive because

$$980\alpha^3 + 552\alpha^2 - 4257\alpha - 4207 = -(980\alpha + 1532)(2 - \alpha)(\alpha + 1) - (1143 + 765\alpha)$$

is negative. Using (16) we obtain that $a_0 = 9P_{10}(\alpha)/32a_2(\alpha + 4)^4$ where

$$\begin{aligned} P_{10}(\alpha) = & -20792743232\alpha^{10} - 168248775872\alpha^9 + 54551858544\alpha^8 \\ & + 3066837388032\alpha^7 + 4844633801556\alpha^6 - 14799467270700\alpha^5 - 43681339670379\alpha^4 \\ & - 4381425810042\alpha^3 + 94728169651149\alpha^2 + 113143847999692\alpha + 41092635382468. \end{aligned}$$

Hence, to conclude the proof we need to show that $a_0 \geq 0$ or equivalently that $P_{10}(\alpha) \geq 0$ whenever $-1 \leq \alpha \leq 2$. This follows from the following identity

$$\begin{aligned} P_{10}(\alpha) = & (20792743232\alpha^2 + 189041519104\alpha + 300831606416)(2 - \alpha)^3(\alpha + 1)^5 \\ & + (1295259115248\alpha + 3335882569236)(2 - \alpha)^2(\alpha + 1)^4 \\ & + (1471491213228\alpha + 7953881034231)(2 - \alpha)(\alpha + 1)^3 \\ & + 1343948812407(2 - \alpha)(\alpha + 1)^2 \\ & + (4661891728632\alpha^2 + 11252369540556\alpha + 6746792560920), \end{aligned} \quad (17)$$

since after examining the signs of the different factors it is possible to conclude that each term in the sum is nonnegative. $\square$

REMARK. Checking that this last identity holds constitutes a formal proof of the fact that $P_{10}(\alpha) \geq 0$ for every $-1 \leq \alpha \leq 2$. Explaining how we obtain it is a bit harder, specially since it involves a "trial and error" process. Essentially, we try to divide $P_{10}(\alpha)$ by a polynomial $A(\alpha)$ of degree a which was known to be positive for the desired range. Hence we obtain that $P_{10}(\alpha) = Q(\alpha)A(\alpha) + R(\alpha)$, where the degrees of Q and R are at most $(10 - a)$ and $(a - 1)$. A sufficient condition for $P_{10}(\alpha) \geq 0$ is then that both $Q(\alpha)$ and $R(\alpha)$ are nonnegative for the desired range. Since we have $-1 \leq \alpha \leq 2$, natural candidates for $A(\alpha)$ took the form $(2 - \alpha)^m(\alpha + 1)^n$. The polynomial division can be made easily using a symbolic manipulation package (cf. the function `quo` in MAPLE), while a plotting routine can make an initial assessment of whether the decomposition was successful, i.e. whether both Q and R are nonnegative (if not, we would try again with different m and n). For instance, the first successful division made to arrive to (17) had $A(\alpha) = (2 - \alpha)^3(1 + \alpha)^5$. In a sense this means that we change a degree 10 problem (showing that P_{10} is nonnegative) by two problems having degrees 2 and 7 (showing respectively that Q and R are nonnegative). The same procedure can be repeated for Q and for R and then for their respective quotients and rests and so on until all polynomials involved are either of the form $(2 - \alpha)^m(\alpha + 1)^n$ or have at most degree 2 and their roots (hence their signs) can be obtained analytically. After doing all these divisions it is easy to put back everything together into a unique decomposition as in (17). Of course, we have no guarantee that this procedure would work for any polynomial, but it worked for P_{10} .

Acknowledgements

This research was partially funded by a CAPES-PROCAD grant. The author is also grateful to the Department of Mathematics of the Università di Roma "La Sapienza" and specially to Prof. F. Spizzichino for support during a sabbatical leave.

References

- [1] V. A. Volkonskij and J. A. Rozanov, "Some limit theorems for random functions - I," *Theory Prob. Appl.*, vol. 4, pp. 178 – 197, 1959.
- [2] A. Fedotov, P. Harremoës, and F. Topsøe, "Refinements of Pinsker's Inequality," *IEEE Trans. Inf. Theory*, vol. 49, pp. 1491–1498, June 2003.
- [3] A. R. Barron, "Entropy and the Central Limit Theorem," *Annals Probab.*, vol. 14, no. 1, pp. 336 – 342, 1986.
- [4] F. Topsøe, "Information theoretical optimization techniques," *Kybernetika*, vol. 15, no. 1, pp. 8 – 27, 1979.
- [5] P. Harremoës and P. Ruzankin, "Rate of convergence to Poisson law in terms of information divergence," *IEEE Trans. Inf. Theory*, vol. 50, no. 9, pp. 2145–2149, 2004.

- [6] M. Del Pino and J. Dolbeault, “Best constants for Gagliardo-Nirenberg inequalities and applications to nonlinear diffusions,” *Journal Mathematiques Pures et Appliquees*, vol. 81, pp. 847–875, 2002.
- [7] I. Csiszár, “Information-type measures of difference of probability distributions and indirect observations,” *Studia Sci. Math. Hungar.*, vol. 2, pp. 299–318, 1967.
- [8] J. H. B. Kemperman, “On the optimal rate of transmitting information,” *Ann. Math. Statist.*, vol. 40, pp. 2156 – 2177, Dec. 1969.
- [9] M. S. Pinsker, *Information and Information Stability of Random Variables and Processes*. A. Feinstein, tr. and ed., San Francisco: Holden-Day, 1964.
- [10] H. P. McKean Jr., “Speed of approach to equilibrium for Kac’s caricature of a Maxwellian gas,” *Arch. Rational Mech. Anal.*, vol. 21, pp. 343 – 367, 1966.
- [11] I. Csiszár, “A note on Jensen’s inequality,” *Studia Sci. Math. Hungar.*, vol. 1, pp. 185–188, 1966.
- [12] S. Kullback, “A lower bound for discrimination information in terms of variation,” *IEEE Trans. Inf. Theory*, vol. IT-13, pp. 126 – 127, Jan. 1967.
- [13] S. Kullback, “Correction to ”a lower bound for discrimination information in terms of variation”,” *IEEE Trans. Inf. Theory*, vol. IT-16, p. 652, 1970.
- [14] I. Vajda, “Note on discrimination information and variation,” *IEEE Trans. Inf. Theory*, pp. 771–773, Nov. 1970.
- [15] F. Topsøe, “Bounds for entropy and divergence of distributions over a two-element set,” *J. Ineq. Pure Appl. Math.*, vol. 2, Article 25, 2001.
- [16] A. Fedotov, P. Harremoës, and F. Topsøe, “Best Pinsker Bound equals Taylor Polynomial of Degree 49,” *Computational Technologies*, vol. 8, pp. 3–14, 2003.
- [17] S. M. Ali and S. D. Silvey, “A general class of coefficients of divergence of one distribution from another,” *J. Roy. Statist. Soc. Ser B*, vol. 28, pp. 131–142, 1966.
- [18] I. Vajda, *Theory of Statistical Inference and Information*. London: Kluwer Academic Press, 1989.
- [19] I. Taneja, “Generalized relative information and information inequalities,” *J. Ineq. Pure Appl. Math.*, vol. 5(1), Article 21, 2004.
- [20] A. Golan, “Information and entropy econometrics—editor’s view,” *Journal of Econometrics*, vol. 107, pp. 1–15, 2002.
- [21] L. Withers Jr., “Some inequalities relating different measures of divergence between two probability distributions,” *IEEE Trans. Inf. Theory*, vol. 45, no. 5, pp. 1728–1735, 1999.
- [22] I. Csiszár, “Generalized cutoff rates and rényi information measures,” *IEEE Trans. Inf. Theory*, vol. 41, pp. 26–34, Jan. 1995.

- [23] D. Dacunha-Castelle, *Ecole d'Ete de Probabilites de Saint-Flour VII-1977*. Berlin, Heidelberg, New York: Springer, 1978.
- [24] L. Le Cam, *Asymptotic Methods in Statistical Theory*. New York: Springer-Verlag, 1986.
- [25] S. S. Dragomir, V. Gluščević, and C. E. M. Pearce, “Csiszár f -divergence, Ostrowski’s inequality and mutual information,” *Nonlinear Analysis*, vol. 47, pp. 2375–2386, 2001.
- [26] F. Topsøe, “Some inequalities for information divergence and related measures of discrimination,” *IEEE Trans. Inf. Theory*, vol. 46, pp. 1602–1609, 2000.
- [27] I. Csiszár, “On topological properties of f -divergences,” *Studia Sci. Math. Hungar.*, vol. 2, pp. 329–339, 1967.
- [28] G. L. Gilardoni, “Accuracy of posterior approximations via chi-squared and harmonic divergences,” *J. of Statistical Planning and Inference*, vol. 128, no. 2, pp. 475–487, 2005.
- [29] G. L. Gilardoni, “Very accurate posterior approximations based on finite mixtures of the hyperparameters conditionals,” *Computational Statistics and Data Analysis*, 2006 (to appear).
- [30] C. Kraft, “Some conditions for consistency and uniform consistency of statistical procedures,” in *Univ. of California Publ. in Statistics*, vol. 1, pp. 125–142, Berkeley: Univ. of California, 1955.
- [31] E. Ordentlich and M. J. Weinberger, “A distribution dependent refinement of Pinsker’s inequality,” *IEEE Trans. Inf. Theory*, vol. 51, pp. 1836–1840, 2005.
- [32] F. Topsøe, “Some bounds for the logarithmic function,” in *RGMIA Research Report Collection*, vol. 7(2), Article 6, 2004. To appear in *Inequalities Theory and Applications*, Vol. 4, Y. J. Cho, J. K. Kim and S. S. Dragomir, eds.